CONTENTS

VOLUME 121 • NUMBER 1 • JANUARY 2003

1 Olaf Andersen. Editorial

3 Roberto Araya, José L. Liberona, J. César Cárdenas, Nora Riveros, Manuel Estrada, Jeanne A. Powell, M. Angélica Carrasco, and Enrique Jaimovich. Dihydropyridine receptors as voltage sensors for a depolarization-evoked, IP_3R-mediated, slow calcium signal in skeletal muscle cells

17 Paola Vergani, Angus C. Nairn, and David C. Gadsby. On the mechanism of MgATP-dependent gating of CFTR Cl^- channels

37 Tamara Rosenbaum, León D. Islas, Anne E. Carlson, and Sharona E. Gordon. Dequalinium: A novel, high-affinity blocker of CNGA1 channels

49 Mahealani K. Monteilh-Zoller, Meredith C. Hermosura, Monica J.S. Nadler, Andrew M. Scharenberg, Reinhold Penner, and Andrea Fleig. TRPM7 provides an ion channel mechanism for cellular entry of trace metal ions

61 Miguel Holmgren. Influence of permeant ions on gating in cyclic nucleotide–gated channels

VOLUME 121 • NUMBER 2 • FEBRUARY 2003

73 Ron R. Kopito and William N. Green. The 56th Annual Meeting and Symposium of the Society of General Physiologists: Trafficking of transporters

81 Karl L. Magleby. *Brief Review:* Gating mechanism of BK (Slo1) channels: So near, yet so far

97 Michiel Helmes, Chee Chew Lim, Ronglih Liao, Ajit Bharti, Lei Cui, and Douglas B. Sawyer. Titin determines the Frank-Starling relation in early diastole

111 Joseph F. Consiglio, Payam Andalib, and Stephen J. Korn. Influence of pore residues on permeation properties in the Kv2.1 potassium channel. Evidence for a selective functional interaction of K^+ with the outer vestibule

125 Xiao-Ming Xia, J.P. Ding, and Christopher J. Lingle. Inactivation of BK channels by the NH_2 terminus of the β2 auxiliary subunit: An essential role of a terminal peptide segment of three hydrophobic residues

149 Michael Duffield, Grigori Rychkov, Allan Bretag, and Michael Roberts. Involvement of helices at the dimer interface in ClC-1 common gating

163 Michael F. Sheets and Dorothy A. Hanck. Molecular action of lidocaine on the voltage sensors of sodium channels

177 Acknowledgement

179 Correction

VOLUME 121 • NUMBER 3 • MARCH 2003

181 Lena Eliasson, Xiaosong Ma, Erik Renström, Sebastian Barg, Per-Olof Berggren, Juris Galvanovskis, Jesper Gromada, Xingjun Jing, Ingmar Lundquist, Albert Salehi, Sabine Sewing, and Patrik Rorsman. SUR1 regulates PKA-independent cAMP-induced granule priming in mouse pancreatic B-cells

199 Huajun Liu, Joshua Atkins, and Robert S. Kass. Common molecular determinants of flecainide and lidocaine block of heart Na^+ channels: Evidence from experiments with neutral and quaternary flecainide analogues

215 Shetuan Zhang, Harley T. Kurata, Steven J. Kehl, and David Fedida. Rapid induction of P/C-type inactivation is the mechanism for acid-induced K^+ current inhibition

227 Andriy Anishkin, Vyacheslav Gendel, Neda A. Sharifi, Chien-Sung Chiang, Lena Shirinian, H. Robert Guy, and Sergei Sukharev. On the conformation of the COOH-terminal domain of the large mechanosensitive channel MscL

245 Thomas Voets, Annelies Janssens, Jean Prenen, Guy Droogmans, and Bernd Nilius. Mg^{2+}-dependent gating and strong inward rectification of the cation channel TRPV6

VOLUME 121 • NUMBER 4 • APRIL 2003

261 Donglin Guo, Yajamana Ramu, Angela M. Klem, and Zhe Lu. Mechanism of rectification in inward-rectifier K^+ channels

277 Guillermo Avila, Kristen M. S. O'Connell, and Robert T. Dirksen. The pore region of the skeletal muscle ryanodine receptor is a primary locus for excitation-contraction uncoupling in central core disease

287 Stéphane Lourdel, Marc Paulais, Pedro Marvao, Antoine Nissant, and Jacques Teulon. A chloride channel at the basolateral membrane of the distal-convoluted tubule: A candidate ClC-K channel

301 Mia Löfgren, Eva Ekblad, Ingo Morano, and Anders Arner. Nonmuscle myosin motor of smooth muscle

311 W.K. Chandler, S. Hollingworth, and S.M. Baylor. Simulation of calcium sparks in cut skeletal muscle fibers of the frog

325 Tong Mook Kang, Vladislav S. Markin, and Donald W. Hilgemann. Ion fluxes in giant excised cardiac membrane patches detected and quantified with ion-selective microelectrodes

VOLUME 121 • NUMBER 5 • MAY 2003

349 C. William Balke and L. Goldman. *Commentary:* Excitation contraction coupling in cardiac muscle: Is there a purely voltage-dependent component?

353 H. Griffiths and K.T. MacLeod. The voltage-sensitive release mechanism of excitation contraction coupling in rabbit cardiac muscle is explained by calcium-induced calcium release

375 Jin Bong Park, Hee Jeong Kim, Pan Dong Ryu, and Edward Moczydlowski. Effect of phosphatidylserine on unitary conductance and Ba^{2+} block of the BK Ca^{2+}–activated K^+ channel: Re-examination of the surface charge hypothesis

399 Jorge Ramos, Wonyong Jung, Josefina Ramos-Franco, Gregory A. Mignery, and Michael Fill. Single channel function of inositol 1,4,5-trisphosphate receptor type-1 and -2 isoform domain-swap chimeras

413 Warren G. Hill, Marcia A. Kaetzel, Bellamkonda K. Kishore, John R. Dedman, and Mark L. Zeidel. Annexin A4 reduces water and proton permeability of model membranes but does not alter aquaporin 2–mediated water transport in isolated endosomes

427 Richard W. Carr, Svetlana Pianova, Juana Fernandez, James B. Fallon, Carlos Belmonte, and James A. Brock. Effects of heating and cooling on nerve terminal impulses recorded from cold-sensitive receptors in the guinea-pig cornea

441 Alan Fein. Inositol 1,4,5-trisphosphate–induced calcium release is necessary for generating the entire light response of *Limulus* ventral photoreceptors

451 Jürgen Rettinger and Günther Schmalzing. Activation and desensitization of the recombinant $P2X_1$ receptor at nanomolar ATP concentrations

463 Tatyana I. Rokitskaya, Elena A. Kotova, and Yuri N. Antonenko. Tandem gramicidin channels cross-linked by streptavidin

477 Robyn L. Goforth, Aung K. Chi, Denise V. Greathouse, Lyndon L. Providence, Roger E. Koeppe II, and Olaf S. Andersen. Hydrophobic coupling of lipid bilayer energetics to channel function

VOLUME 121 • NUMBER 6 • JUNE 2003

495 Heather L. Agler, Jenafer Evans, Henry M. Colecraft, and David T. Yue. Custom distinctions in the interaction of G-protein β subunits with N-type ($Ca_V2.2$) versus P/Q-type ($Ca_V2.1$) calcium channels

511 Karel Talavera, Annelies Janssens, Norbert Klugbauer, Guy Droogmans, and Bernd Nilius. Extracellular Ca^{2+} modulates the effects of protons on gating and conduction properties of the T-type Ca^{2+} channel α_{1G} ($Ca_V3.1$)

529 Karel Talavera, Annelies Janssens, Norbert Klugbauer, Guy Droogmans, and Bernd Nilius. Pore structure influences gating properties of the T-type Ca^{2+} channel α_{1G}

541 Valérie Leuranguer, Robert T. Dirksen, and Kurt G. Beam. Potentiated L-type Ca^{2+} channels rectify

551 Bhavna Tanna, William Welch, Luc Ruest, John L. Sutko, and Alan J. Williams. An anionic ryanoid, 10-O-succinoylryanodol, provides insights into the mechanisms governing the interaction of ryanoids and the subsequent altered function of ryanodine-receptor channels

563 Galen E. Flynn and William N. Zagotta. A cysteine scan of the inner vestibule of cyclic nucleotide–gated channels reveals architecture and rearrangement of the pore

583 Jixin Wang, Joseph J. Salata, and Paul B. Bennett. Saxitoxin is a gating modifier of hERG K^+ channels

599 Jie Liu, Mei Zhang, Min Jiang, and Gea-Ny Tseng. Negative charges in the transmembrane domains of the HERG K channel are involved in the activation- and deactivation-gating processes

615 Vladimir V. Cherny, Ricardo Murphy, Valerij Sokolov, Richard A. Levis, and Thomas E. DeCoursey. Properties of single voltage-gated proton channels in human eosinophils estimated by noise analysis and by direct measurement